

ORIGINAL ARTICLE

Knowledge and perception of patients towards the hazards and dose of diagnostic radiation in Bahrain

Khalid Hameed Alshammari^{1*}, Layla Emad Alramadan¹, Jaffar Mohamed Hasan Ali¹, Musaab Jamal Farraj¹, Khaled Ab dulaziz Alanazi¹, Alaa Waleed Ahmed Almurbati¹

ABSTRACT

Background: Ionizing radiation is a helpful diagnostic tool in a wide range of medical specialties. However, it imposes a risk of radiation exposure to both patients and healthcare professionals. Therefore, this study aimed at evaluating the level of knowledge of patients toward radiation exposure hazards and radiation dose.

Methodology: The study was a qualitative cross-sectional study. Data were collected through a self-administered online questionnaire that was distributed electronically to the patients who underwent the diagnostic radiological procedure. Data were represented in terms of frequencies and valid percentages for categorical variables. A one-way analysis of variance test was used to compare numerical variables between subgroups.

Results: The mean knowledge score was below average (5.08 ± 2.952). Patients aged between 18 and 25 years, and single patients had significantly higher mean scores. Patients who were advised about the hazards of radiation showed a significantly higher mean score. These informed patients had the highest mean score in the entire sample population.

Conclusion: The level of knowledge of patients in Bahrain toward risks of radiation exposure is inadequate. Further research is required on a national level. Awareness campaigns are highly recommended to improve the level of knowledge.

Keywords: Knowledge, perception of patients, the hazards, dose of diagnostic radiation, Bahrain.

Introduction

Ionizing radiation is considered a non-invasive diagnostic intervention [1], which can help in the process of decision making regarding a medical condition in terms of pharmacological or interventional strategy [2]. The ionizing radiation is used in a couple of techniques, including X-ray, computed tomography (CT) scan, and magnetic resonance imaging (MRI) [3].

Individuals who are at the highest exposure level of ionizing radiation during the intervention, either for treatment or diagnostic purposes, are healthcare professionals and the patients [4]. The radiation doses can be simply estimated by a dosimeter [5]. However, the dose of radiation exposure to patients or medical staff cannot be directly observed with naked eyes while measuring by a dosimeter [6].

Although healthcare professionals working in radiological specialties have good training in radiation safety [7], patient's education is also necessary in order

to minimize their risk of cancer or any other hazards of radiation exposure [8].

Medical professionals, including vascular surgeons, cardiologists, and gastroenterologists, have training on the hazards of radiation exposure [9]. This is of particular importance in the case of intravascular interventions [10]. These procedures impose a similar risk of radiation exposure on both patients and doctors [11]. Therefore, monitoring the radiation exposure is mandatory [12].

Correspondence to: Khalid Hameed Alshammari

*Intern Doctor, Salmaniya Medical Complex, Manama, Bahrain.

Email: khalidalshammari78@gmail.com

Full list of author information is available at the end of the article.

Received: 22 October 2019 | **Accepted:** 17 November 2019

Other types of interventions may require a higher radiation dose, which is mainly required for some patient's clinical factors and certain amount of dose is required for diagnosis [13]. The medical staff can be protected by using dosimeter, which determines the degree of exposure [14]. Previous reports showed that patients receive a significantly higher dose of radiation during diagnostic procedures [15].

Several studies have explored the level of knowledge of healthcare professionals toward diagnostic radiation hazards [16] though there is scarcity of data about the level of knowledge of patients about the risk of ionizing radiation that they are exposed to Alzahrani et al. [17]. Therefore, the aim of this study was to explore the level of knowledge of the Bahrain population toward the risks and hazards of radiation used for diagnostic procedures and radiation dose.

Subjects and Methods

A qualitative, cross-sectional study was performed in Manama, Bahrain, where an online self-developed questionnaire was distributed, via a link to Google forms to the patients who underwent the diagnostic radiological procedure. Only those who completed the questionnaire were included in the analysis.

Data were collected through a self-administered questionnaire. The responses were divided into two sections: the first section included questions on socio-demographic data. The second section was questioned on knowledge of radiation exposure hazards and radiation doses in different diagnostic procedures.

The knowledge of respondents was evaluated by calculating the scores for correct answers. Each correct response was given one point. Data were represented in terms of frequencies and valid percentages for categorical variables. Mean, standard deviations, minimum, and maximum values were used to describe the numerical variables. A one-way analysis of variance (ANOVA) test was used to compare numerical variables between the subgroups. All p -values <0.05 were considered as statistically significant. IBM Statistical Package for the Social Science (IBM Corp, Armonk, NY) was used to perform all statistical calculations, version 21 for Microsoft Windows.

Results

Three hundred and fifty-seven participants responded to this online questionnaire in this study. Out of 357 participants, age was subcategorized into four groups, starting with 18–25 years old and ending with more than 56-year old. Most of the respondents (58%) belonged to the age group 18–25 years old. On the other hand, the age group who were greater than 56-year old had the least number of responses (3.4%).

Single participants constituted 57.7% of participants, while widowed participants were 0.8%. The educational

level was also evaluated, in which 74.8% had a university degree or higher, while 0.8% only were illiterate. All socio-demographic data are shown in detail in Table 1.

Professions of participants were divided into either medical or non-medical professions. Medical professions represented 42.3% of the whole, while only 27.7% had non-medical professions, as shown in Figure 1.

Participants were asked about the type of information provided to them regarding their radiological investigation. It was shown that 65.3% of the patients were told the reason for their investigation, while 63.6% were not informed about the radiation dose that they will be exposed to. Additionally, 73.7% of patients were not told about the hazards of exposure to radiation, and 67.5% were not told about other related risks, as shown in Figure 2.

Participants were also asked about their knowledge about the use of radiation in different diagnostic procedures, where 65.5% of patients knew that ionizing radiation was used in X-ray, 48.2% mentioned that radiation was used in CT, 35% agreed that radiation was used in ultrasound. Also, 41.5% responded with a positive response that MRI required radiation.

On the other hand, 38.4% and 36.4% of patients mentioned that radiation was not used in barium swallowing test and rectal barium test, respectively. All responses are shown in detail in Table 2.

Patients were also asked about their knowledge of exposure to radiation during a diagnostic procedure. The questions revealed that 48.7% mentioned that radiation exposure does not increase the risk of cancer, while 56.3% of participants

Table 1. Socio-demographic data of the respondents.

	Frequency (n)	Percentage (%)
Age group		
18–25	207	58.0
26–40	85	23.8
41–55	43	12.0
>56	12	3.4
Marital status		
Single	206	57.7
Widowed	3	0.8
Married	129	36.1
Divorced	12	3.4
Educational level		
Illiterate	3	0.8
Primary	4	1.1
Secondary	60	16.8
Intermediate	13	3.6
University education and higher	267	74.8




Figure 1. Profession of participants.




Figure 2. Information provided by treating physician.

agreed that repetitive exposure to radiation could increase the cumulative risk of cancer. Also, 37.3% and 42.6% of participants denied being exposed to radiation in airports and at home, respectively. Additionally, 23.5% showed that the risk of cancer due to CT in adults is 1/1,000. Moreover, 91.6% of patients mentioned that they should be told about the reason for X-ray if needed for them. All questions and responses are shown in Table 3.

The total knowledge score was calculated for included questions evaluating knowledge about the dose of radiation and radiation hazards. The minimum score report was zero, while the maximum score was 13. The average knowledge score was 5.08 ± 2.952 . The mean for total score was compared over different demographic

data and influencing factors using one-way ANOVA at the level of significance p -value <0.05 .

It was shown that patients aged between 18 and 25 years old had significantly higher (p -value <0.001) mean score (5.79). Additionally, single patients had significantly higher (p -value = 0.009) mean scores as compared to other groups (5.41).

Patients who were told about the hazards of radiation showed a significantly higher mean score (p -value <0.001) as compared to those who were not provided with this information. It is worth mentioning that this group of patients had the highest mean score as compared to the whole sample population (8.28). All comparisons are shown in Table 4.

Table 2. Knowledge of participants toward radiation use.

		Frequency (n)	Percentage (%)
X-ray	I don't know	21	5.9
	No	28	7.8
	Yes	234	65.5
CT	I don't know	24	6.7
	No	63	17.6
	Yes	172	48.2
Ultrasound	I don't know	27	7.6
	No	95	26.6
	Yes	125	35.0
MRI	I don't know	24	6.7
	No	89	24.9
	Yes	148	41.5
Barium swallowing test	I don't know	45	12.6
	No	137	38.4
	Yes	58	16.2
Barium rectal test	I don't know	45	12.6
	No	130	36.4
	Yes	59	16.5

Table 3. Knowledge of patients towards radiation dose and risk of radiation exposure.

		Frequency (n)	Percentage (%)
Radiation can increase the risk of cancer	I don't know	30	8.4
	No	174	48.7
	Yes	75	21.0
The cumulative risk of cancer results from repetitive exposure to radiation	I don't know	30	8.4
	No	51	14.3
	Yes	201	56.3
I am exposed to radiation in Airports	I don't know	30	8.4
	No	133	37.3
	Yes	115	32.2
I am exposed to radiation at home	I don't know	33	9.2
	No	152	42.6
	Yes	101	28.3
Have you heard before about posteroanterior radiation	I don't know	18	5.0
	No	264	73.9
	Yes	16	4.5
The total amount of radiation received from posteroanterior radiation is lower than the amount produced from chest X-ray	I don't know	48	13.4
	No	107	30.0
	Yes	101	28.3
The risk of cancer due to CT radiation in adults is	1/1,000	84	23.5
	1/2,000	51	14.3
	1/3,000	64	17.9
	1/4,000	7	2.0
	1/5,000	59	16.5
Do you think that you should be told about your need for X-ray?	No	10	2.8
	Yes	327	91.6

Table 4. Comparison of total knowledge score over different socio-demographic data.

		Mean knowledge score	p-value
Age group	18–25	5.79	<0.001*
	26–40	4.26	
	41–55	4.28	
	>56	2.75	
Marital status	Single	5.41	0.009*
	Widowed	5.00	
	Married	4.77	
	Divorced	2.75	
Educational level	Illiterate	3.00	0.143
	Primary	5.00	
	Secondary	5.00	
	Intermediate	5.26	
	University education and higher	3.38	
Profession	Medical	5.55	0.380
	None medical	5.23	
Did your doctor mention the hazards of radiation	No	5.10	<0.001*
	Yes	8.28	

*p-value at the level of significance <0.05.

Discussion

Ionizing radiation played a very important role in the advancement of radiological investigations over the past decade [8]. However, some adverse effects can occur due to its use although the occurrence of these events is very rare [6]. Knowledge of patients about these hazards is crucial [9].

In the present study, the level of knowledge of Bahrain patients toward radiation dose and risk of exposure to radiation during diagnostic procedures was evaluated. It was revealed that the level of knowledge of patients was below average, where the mean knowledge score was 5.08 ± 2.952 . It was also shown that patients aged between 18 and 25 years old, and single patients had significantly higher mean score. Also, patients who were told the hazards of radiation showed a significantly higher mean score.

The level of patients' knowledge was also evaluated in different clinical settings. Almatared et al. [18] investigated the level of knowledge of patients toward ionizing radiation on 375 patients and showed that the level of knowledge of patients was low in Nijran city, Saudi Arabia. Similarly, the present study showed that the mean score for the knowledge section was below average. Additionally, the present study revealed that patients who got advice from their doctors about the hazards of exposure to ionizing radiation had the highest mean in the entire sample population.

Moreover, Sin et al. [19] evaluated the awareness of the patients towards risks of radiological diagnostic

procedures. The study recruited 173 patients only who underwent either CT or X-ray imaging. Sin et al. [19] revealed that the awareness of patients toward radiation safety and the risk of exposure were unsatisfactory. Additionally, Sin et al. [19] recommended awareness programs to improve the level of knowledge of patients towards radiation exposure risk.

Findings of Sin et al. [19] were similar to the findings of the present work; however, the present work recruited a larger sample size, which increases the reliability of the present outcomes. Moreover, the present work demonstrated the important role of physicians in increasing the knowledge and perception of patients towards radiation exposure risks.

Takakuwa et al. [20] evaluated the level of knowledge of patients toward risk of CT investigations, specifically in the Emergency Department. Takakuwa et al. [20] recruited 383 patients who were admitted to the Emergency Department and were required to undergo a CT investigation for diagnostic purpose, and showed that the patients were not aware of the extent of risk due to exposure to radiation, especially the risk of cancer. Furthermore, Takakuwa et al. [20] showed that age, education level, and race were different factors influencing the level of knowledge of patients, as well as their attitude towards radiation exposure risk.

The present study also described a significant difference in the level of knowledge between different age groups, where the age group of 18–25 years showed a significantly higher score as compared to other age groups. However, the present work could not find a

significant difference in the level of knowledge among different educational levels.

The present study had some limitations; it was performed in only one area in Bahrain, multicenter studies in the future are recommended. Furthermore, the small sample size could decrease the reliability of the results.

Conclusion

The level of knowledge of Bahrain patients toward the risk of exposure to radiation and radiation dose is relatively low. National awareness programs should be encouraged to increase their level of knowledge regarding the risk of this exposure. Additionally, healthcare professionals should be encouraged to advise their patients about the procedure that they will have its hazards and methods of prevention.

List of Abbreviations

CT Computed tomography
MRI Magnetic resonance imaging

Conflict of interest

The authors declare that there is no conflict of interest regarding the publication of this article.

Funding

None.

Consent for publication

Informed consent was obtained from all participants.

Ethical approval

A statement was included at the beginning of the questionnaire clarifying that the participation in the study is totally voluntary and that collected data were anonymous and only used for the purpose of the study.

Author details

Khalid Hameed Alshammari¹, Layla Emad Alramadan¹, Jaffar Mohamed Hasan Ali¹, Musaab Jamal Farraj¹, Khaled Abdulaziz Alanazi¹, Alaa Waleed ahmed almurbati¹
1. Intern Doctor, Salmaniya Medical Complex, Manama, Bahrain

References

- Ahmed RM, Elamin AM, Elsamani M, Hassan WB. Knowledge and performance of radiographers towards radiation protection, Taif, Saudi Arabia. *IOSR J Dent Med Sci*. 2015;14(3):63–8.
- Al Mohiy H, Sim J, Seeram E, Davidson R. Differences between Saudi Arabian and Australian radiographers' knowledge and attitudes about paediatric CT doses. *Radiographer*. 2011;58(3):21–7. <https://doi.org/10.1002/j.2051-3909.2011.tb00151.x>
- Hagi S, Khafaji M. Medical student's knowledge of ionizing radiation and radiation protection. *Saudi Med J*. 2011;32(5):520–4.
- Al Dossari HM, Al Dosary A, Al Rashed A, Alamro S, Al qahtani F, Alanazi A, et al. Assessing the level of knowledge about radiation dose in common radiological examinations among physicians in Riyadh. *Int J Pharm Res Allied Sci*. 2018;7(1):5.
- Arnout EA, Jafar A. Awareness of biological hazards and radiation protection techniques of dental imaging-a questionnaire based cross-sectional study among Saudi dental students. *J Dent Health Oral Disord Ther*. 2014;1(1):1–7. <https://doi.org/10.15406/jdhodt.2014.01.00008>
- Enabulele JE, Itimi E. Endodontic radiology, practice, and knowledge of radiation biology, hazard, and protection among clinical dental students and interns. *Saudi Endod J*. 2015;1(3):171. <https://doi.org/10.4103/1658-5984.163634>
- Al Mohiy H, Sim J, Seeram E, Annabell N, Geso M, Mandarano G, et al. A dose comparison survey in CT departments of dedicated paediatric hospitals in Australia and Saudi Arabia. *World J Radiol*. 2012;4(10):431. <https://doi.org/10.4329/wjrv.v4.i10.431>
- Al-Rammah TY. CT radiation dose awareness among pediatricians. *Ital J pediatr*. 2016;42(1):77. <https://doi.org/10.1186/s13052-016-0290-3>
- Almohiy H. Multicenter study assessing ophthalmologist's knowledge towards radiation dose when prescribing CT scans for pediatric patients: a Saudi Arabian perspective. *J Radiat Res Appl Sci*. 2016;9(1):96–100. <https://doi.org/10.1016/j.jrras.2015.10.001>
- Almaghrabi N. A cross-sectional study of knowledge and awareness of radiation exposure risk in Makkah, Saudi Arabia. *Am J Res Commun*. 2016;4(10):67–75. <https://doi.org/10.21474/IJAR01/1008>
- Alotaibi AT, Alnafea MA. Radiation dose awareness among radiology staff. *Austin J Nucl Med Radiother*. 2017;4(1):1023.
- ElOlemy AT, Almuwannis MS, Alamiri MS, Alkhudair KM. Knowledge attitude and practices of ultrasound safety in pregnancy among healthcare providers in Riyadh, Saudi Arabia: a survey study. *Tanta Med J*. 2015;43(3):98. <https://doi.org/10.4103/1110-1415.162445>
- Saeed MK, Al-shaari H, Almarzooq MM, Alsareii SA, Aljerdah SA, Al-ayed MS. Radiation awareness among physicians about the hazards of radiological examinations on the health of workers and their patients in Saudi Arabia. *J Radiat Res Appl Sci*. 2018;11(4):299–304. <https://doi.org/10.1016/j.jrras.2018.04.001>
- Taha MT, Al-Ghorabie FH, Kutbi RA, Saib WK. Assessment of entrance skin doses for patients undergoing diagnostic X-ray examinations in King Abdullah Medical City, Makkah, KSA. *J Radiat Res Appl Sci*. 2015;8(1):100–3. <https://doi.org/10.1016/j.jrras.2014.12.003>
- Al Mohiy H, Sim J, Annabell N, Seeram E, Davidson R. A comparison of Saudi Arabian and Australian radiographers' perceptions of computed tomography radiation dose. *Radiographer*. 2012;59(4):113–8. <https://doi.org/10.1002/j.2051-3909.2012.tb00186.x>
- Rahmatullah TA, Alela KA, Alanazi KS. Medical student's knowledge of ionizing radiation and radiation protection in Riyadh, Saudi Arabia. *Egypt J Hosp Med*. 2018;70(1). <https://doi.org/10.12816/0042969>
- Alzahrani YA, Alomairi NE, Abdelbaky A, Althagafi R, Altalhi M, Althagafi M, et al. Awareness of medical staff towards radiation protection and dose for radiological procedures in Taif, Saudi Arabia. *Int J Med Sci Clin Invent*. 2018;5(5):3827–32.

18. Almatared MG, Almansour AM, Alquidaihi AE, Agbaje HO, Almutared AS. Knowledge and Attitude towards Ionizing Radiation among Patients Attending the King Khalid Hospital Najran, Saudi Arabia. *J Biosci Med.* 2017;5(10):75. <https://doi.org/10.4236/jbm.2017.510008>
19. Sin HK, Wong CS, Huang B, Yiu KL, Wong WL, Chu YC. Assessing local patients' knowledge and awareness of radiation dose and risks associated with medical imaging: a questionnaire study. *J Med Imaging Radiat Oncol.* 2013;57(1):38–44. <https://doi.org/10.1111/j.1754-9485.2012.02471.x>
20. Takakuwa KM, Estepa AT, Shofer FS. Knowledge and attitudes of emergency department patients regarding radiation risk of CT: effects of age, sex, race, education, insurance, body mass index, pain, and seriousness of illness. *Am J Roentgenol.* 2010;195(5):1151–8. <https://doi.org/10.2214/AJR.09.3847>